

01	A	B	C	D	E
02	A	B	C	D	E
03	A	B	C	D	E
04	A	B	C	D	E
05	A	B	C	D	E
06	A	B	C	D	E
07	A	B	C	D	E
08	A	B	C	D	E
09	A	B	C	D	E
10	A	B	C	D	E
11	A	B	C	D	E
12	A	B	C	D	E
13	A	B	C	D	E

14	A	B	C	D	E
15	A	B	C	D	E
16	A	B	C	D	E
17	A	B	C	D	E
18	A	B	C	D	E
19	A	B	C	D	E
20	A	B	C	D	E
21	A	B	C	D	E
22	A	B	C	D	E
23	A	B	C	D	E
24	A	B	C	D	E
25	A	B	C	D	E
26	A	B	C	D	E

27	A	B	C	D	E
28	A	B	C	D	E
29	A	B	C	D	E
30	A	B	C	D	E
31	A	B	C	D	E
32	A	B	C	D	E
33	A	B	C	D	E
34	A	B	C	D	E
35	A	B	C	D	E
36	A	B	C	D	E
37	A	B	C	D	E
38	A	B	C	D	E
39	A	B	C	D	E

PRACTICE QUESTIONS

TTBU - The Testing Board Ukraine

2PHRU

Krok 2 Pharmacy (Russian)

EDITION

Demo

QUESTIONS

9

ISSUED

September 3, 2026

Before you begin

What this is

9 practice questions for TTBU - The Testing Board Ukraine **2PHRU**, each with its answer key and an explanation. Answer a question before you read its key — the explanation is worth more once you have committed.

If something looks wrong

Send us three things and we will check it:

- the exam code, **2PHRU**
- the question number
- the email address on your order

Write to **support@mometrix.net**. We reply within one business day.

Licence

This file is licensed to one person. Sharing or republishing it, in whole or in part, ends that licence.

QUESTION 1

SINGLE CHOICE

Which non-steroidal anti-inflammatory drug (NSAID) is a selective inhibitor of cyclooxygenase-2 (COX-2)?

- ☐ A Ibuprofen
- ☐ B Diclofenac
- ☒ C Celecoxib
- ☐ D Aspirin

ANSWER C

WHY Celecoxib is a selective COX-2 inhibitor used for its anti-inflammatory effect with reduced gastrointestinal toxicity compared with non-selective NSAIDs. Ibuprofen, diclofenac and aspirin are non-selective COX inhibitors.

QUESTION 2

SINGLE CHOICE

Which group of bronchodilators acts primarily through stimulation of β_2 -adrenergic receptors of the bronchi?

- ☐ A M-cholinoblockers
- ☒ B β_2 -adrenomimetics
- ☐ C Xanthine derivatives
- ☐ D Leukotriene receptor antagonists

ANSWER B

WHY β_2 -adrenomimetics (e.g., salbutamol, fenoterol) produce bronchodilation by activating β_2 -adrenergic receptors on bronchial smooth muscle. M-cholinoblockers (e.g., ipratropium bromide) act via blockade of muscarinic receptors, xanthines (e.g., theophylline) inhibit phosphodiesterase, and leukotriene receptor antagonists (e.g., montelukast) block the cysteinyl leukotriene receptor.

QUESTION 3

SINGLE CHOICE

To which pharmacological group does the antibiotic amoxicillin belong?

- ☐ A Macrolides
- ☐ B Aminoglycosides
- ☐ C Fluoroquinolones
- ☒ D Semisynthetic penicillins

ANSWER D

WHY Amoxicillin is a semisynthetic penicillin of the β -lactam class with a broad spectrum of activity, especially against Gram-negative organisms. Macrolides (e.g., azithromycin), aminoglycosides (e.g., gentamicin) and fluoroquinolones (e.g., ciprofloxacin) are different antibacterial classes.

QUESTION 4

SINGLE CHOICE

Which medicinal plant is the main natural source of the alkaloid morphine?

- ☐ A Atropa belladonna
- ☒ B Papaver somniferum
- ☐ C Vinca minor
- ☐ D Rauwolfia serpentina

ANSWER B

WHY Morphine is the principal alkaloid of the opium poppy (Papaver somniferum). Atropa belladonna contains tropane alkaloids (atropine, scopolamine), Vinca minor contains vincamine, and Rauwolfia serpentina contains reserpine.

QUESTION 5

SINGLE CHOICE

A pharmacist is preparing an injectable solution of a drug that is poorly soluble in water. Which of the following excipients would be MOST appropriate to increase aqueous solubility?

- ☐ A Liquid paraffin
- ☐ B Sodium chloride
- ☒ C Polyethylene glycol 400 (PEG 400)
- ☐ D Magnesium stearate

ANSWER C

WHY PEG 400 is a water-miscible cosolvent widely used in pharmaceutical technology to increase the aqueous solubility of poorly water-soluble drugs. Liquid paraffin is an oily vehicle, sodium chloride is mainly used for isotonicity adjustment, and magnesium stearate is a hydrophobic lubricant used in tablet manufacture.

QUESTION 6

SINGLE CHOICE

Omeprazole belongs to which pharmacological group of antiulcer drugs?

- ☐ A H₂-histamine receptor blockers
- ☒ B Proton pump inhibitors
- ☐ C Antacids
- ☐ D M-cholinoblockers

ANSWER B

WHY Omeprazole irreversibly inhibits the H⁺/K⁺-ATPase (proton pump) of gastric parietal cells, strongly suppressing gastric acid secretion. H₂-blockers (e.g., ranitidine) block H₂ receptors, antacids neutralize existing acid, and M-cholinoblockers reduce acid via muscarinic blockade.

QUESTION 7 SINGLE CHOICE

A long-term high-dose use of glucocorticoids is most characteristically associated with which adverse effect?

- ☒ **A** Cushing's syndrome (iatrogenic hypercortisolism)
- ☐ **B** Hepatotoxicity with jaundice
- ☐ **C** Severe bradycardia
- ☐ **D** Hypothyroidism

ANSWER A

WHY Prolonged systemic glucocorticoid therapy reproduces the clinical picture of hypercortisolism: moon face, central obesity, striae, hyperglycemia, osteoporosis and immunosuppression. The other listed effects are typical of other drug groups (hepatotoxicity — e.g., methotrexate, isoniazid; bradycardia — β -blockers; hypothyroidism — lithium, amiodarone).

QUESTION 8 SINGLE CHOICE

Diazepam, a widely used anxiolytic, belongs to which pharmacological group?

- ☐ **A** Barbiturates
- ☒ **B** Benzodiazepines
- ☐ **C** Tricyclic antidepressants
- ☐ **D** Atypical antipsychotics

ANSWER B

WHY Diazepam is a long-acting benzodiazepine that potentiates the action of GABA at the GABA-A receptor complex, producing anxiolytic, sedative, muscle relaxant and anticonvulsant effects. Barbiturates, tricyclic antidepressants and atypical antipsychotics have different structures and mechanisms of action.

QUESTION 9

MULTIPLE CHOICE

A patient with chronic heart failure (NYHA class II-III) and reduced left ventricular ejection fraction requires addition of a drug that prolongs survival. Which classes of drugs have been shown to reduce mortality in this patient population? (Choose two.)

- ☒ **A** ACE inhibitors (e.g., enalapril)
- ☐ **B** Short-acting dihydropyridine calcium channel blockers (e.g., nifedipine)
- ☒ **C** Mineralocorticoid receptor antagonists (e.g., spironolactone)
- ☐ **D** Class IC antiarrhythmics (e.g., propafenone)

ANSWER A - C

WHY ACE inhibitors reduce mortality and morbidity in heart failure with reduced ejection fraction. Mineralocorticoid receptor antagonists such as spironolactone and eplerenone also reduce mortality in NYHA II-IV HF (RALES trial). Short-acting dihydropyridines may worsen HF, and Class IC antiarrhythmics are not indicated and may be harmful in structural heart disease.